

Partial permutohedra


Laura Escobar

Washington University in St. Louis

Women at the intersection of mathematics and theoretical physics
meet in Okinawa

March 23, 2023

Based on: arXiv:2207.14253.




Joint with Roger Behrend, Federico Castillo, Anastasia Chavez, Alexander Diaz-Lopez, Pamela Harris and Erik Insko.

Polytopes

The **convex hull** of a set $S \subseteq \mathbb{R}^d$ is the smallest convex set containing S . We denote it by $\text{Conv}(S)$.

Example. The convex hull of $\{(3, 1), (4, 2), (7, 0), (10, 3), (0, 3), (0, 4)\}$ is:



A **polytope** is the convex hull of a finite set.

An invitation to measure polytopes

The geometry of linear optimization is understood by studying polytopes.

The number of k -dimensional regions of a hyperplane arrangement in \mathbb{R}^n is equal to the number of $(n - k)$ -dimensional faces of a zonotope.

Using the rich correspondence between geometry of toric varieties and combinatorics of convex polytopes, Batyrev constructed classes Calabi-Yau manifolds as hypersurfaces of toric varieties and proved the mirror symmetry conjecture for smooth toric varieties.

The n -dimensional associahedron is a polytope whose number of vertices equals the $(n + 1)$ -th Catalan number. Recently, Mizera and Arkani-Hamed–Bai–He–Yan showed that this polytope plays a central role in the theory of scattering amplitudes.




Given a polytope, how many vertices does it have? How many faces?
What's its volume?

Theorem. (Dryer 83, Linial 86, Dryer–Frieze 88) Computing any of these values is $\#P$ -hard.

Partial permutohedra

Definition. (Heuer–Striker '22) Given $d, n \in \mathbb{Z}_{>0}$, the **partial permutohedron** $\mathcal{P}(d, n)$ is the convex hull of all vectors in $\{0, 1, \dots, n\}^d$ whose nonzero entries are distinct.




Example. Let $d = 2$.



Partial permutohedra

Definition. (Heuer–Striker '22) Given $d, n \in \mathbb{Z}_{>0}$, the **partial permutohedron** $\mathcal{P}(d, n)$ is the convex hull of all vectors in $\{0, 1, \dots, n\}^d$ whose nonzero entries are distinct.

Example. Let $d = 3$.



Number of faces

The **f-polynomial** of a d -dimensional polytope \mathcal{P} is

$f_{\mathcal{P}}(t) = \sum_{i=0}^d f_i(\mathcal{P}) t^i$, where $f_i(\mathcal{P})$ denotes the number of i -dimensional faces of \mathcal{P} .

The **Eulerian polynomial** is $A_d(t) = \sum_{i=0}^{d-1} A(d, i) t^i$, where $A(d, i)$ is the number of permutations in S_d with exactly i descents.

A **descent** of w is a position $i < n$ with $w(i) > w(i + 1)$. For example, $w = 3452167$ has descents at positions 3 and 4.

Theorem. (BCCDEHI '22+) The f -polynomial of $\mathcal{P}(d, n)$ with $n \leq d$ is

$$f_{\mathcal{P}(d,n)}(t) = 1 + \sum_{i=0}^{n-1} \sum_{j=1}^{d-i} \binom{d}{i} A_i(t+1) (t+1)^j.$$

Volume of $\mathcal{P}(d, n)$ with $n \geq d - 1$

To simplify volume expressions, we use the normalized volume defined such that $\text{Vol}([0, 1]^m) = m!$.

Denote by $v(d, n)$ the normalized volume of $\mathcal{P}(d, n)$.

Theorem. (BCCDEHI '22+) For any d and n with $n \geq d - 1$, the normalized volume of $\mathcal{P}(d, n)$ is given recursively by

$$v(d, n) = (d - 1)! \sum_{k=1}^d k^{k-2} \frac{v(d - k, n - k)}{(d - k)!} \left(kn - \binom{k}{2} \right) \binom{d}{k},$$

with the initial condition $v(0, n) = 1$.


Examples.

$$\begin{aligned} v(1, n) &= n, \\ v(2, n) &= -1 + 2n^2, \\ v(3, n) &= -6 - 9n + 6n^3, \\ v(4, n) &= -54 - 96n - 72n^2 + 24n^4, \\ v(5, n) &= -840 - 1350n - 1200n^2 - 600n^3 + 120n^5. \end{aligned}$$

Ideas behind the proof

Lemma. Let \mathbf{v} be a vertex of $\mathcal{P}(d, n)$. For each facet \mathcal{F} that does not contain \mathbf{v} , form the pyramid $\text{Pyr}(\mathcal{F}, \mathbf{v})$. The collection of these pyramids for all such facets gives a polyhedral subdivision of $\mathcal{P}(d, n)$, and thus

$$\text{Vol}(\mathcal{P}(d, n)) = \sum_{\substack{\text{facets } \mathcal{F} \\ \mathbf{v} \notin \mathcal{F}}} \text{Vol}(\text{Pyr}(\mathcal{F}, \mathbf{v})).$$



Let

$$\Pi_k = \text{Conv}\{(w(1), \dots, w(k)) \mid w \in S_k\}.$$

Each facet not containing $\mathbf{0}$ is congruent to either a Cartesian product $\Pi_k \times \mathcal{P}(d - k, n - k)$ or Π_d .


This method was used to obtain the volume of $\mathcal{P}(d, n)$ for the case $n = d - 1$ by Amanbayeva–Wang and Strong.

Next, we fix n and vary d .

Volume of $\mathcal{P}(d, 2)$

The following was conjectured in (Heuer–Striker '21).

Theorem. (BCCDEHI '22+) For any d , the normalized volume of $\mathcal{P}(d, 2)$ is $v(d, 2) = 3^d - d$.




$\mathcal{P}(d, 2)$ is equal to $3 \operatorname{Conv}\{\mathbf{0}, \mathbf{e}_1, \dots, \mathbf{e}_d\}$ with d simplices removed.

Volume of $\mathcal{P}(d, 3)$

Theorem. (BCCDEHI '22+) For any d , the normalized volume of $\mathcal{P}(d, 3)$ is

$$v(d, 3) = 6^d - d 3^d - (d - 1) \binom{d}{2}.$$



Theorem. (BCCDEHI '22+) For any d , the normalized volume of $\mathcal{P}(d, 4)$ is

$$v(d, 4) = 10^d - d 6^d - \frac{d(d-1)(d-3)}{6} 3^d - (3d^2 - 6d + 1) \binom{d}{3}.$$

Thank you!